

PREPARING FOR YOUR CATARACT PROCEDURE BEFORE / DURING /AFTER

Name: _____ DOB: _____

You have been scheduled for RIGHT LEFT eye surgery.

DATE OF SURGERY: _____ **Surgeon:** _____

ARRIVAL TIME: Eyesight does not schedule your surgery time. You will be contacted by the surgery center the day before your procedure with your expected arrival time. **You will also be called 1-2 weeks prior to surgery to go over your medical history. If you have not heard from the surgery center by 3:30pm the day before your procedure, please contact them directly at 603-314-8035 to get your time.**

Note: The first 2 pages of this book offer a convenient overview of items covered in this book.

SURGERY CENTER / LOCATION OF SURGERY:

_____ Coastal Surgical Center – 291 Shattuck Way, Newington NH

Alternative locations:

_____ Frisbie Memorial Hospital - 11 Whitehall Road, Rochester NH

_____ Wentworth Douglass Hospital - 789 Central Avenue, Dover NH

_____ Exeter Hospital - 5 Alumni Drive, Exeter NH

AFTER SURGERY APPOINTMENTS: YOU MUST BE SEEN FOR A FOLLOW UP APPOINTMENT AFTER SURGERY. PLEASE PLAN TO SEE US AT EYESIGHT ON THE FOLLOWING DATE/TIME:

1st post-op appointment is in the Portsmouth Office Somersworth Office Exeter Office Kittery, ME Office

on _____ at _____ with Dr. _____.

2nd post-op appointment is in the Portsmouth Office Somersworth Office Exeter Office Kittery, ME Office

on _____ at _____ with Dr. _____.

Light Adjustable / RLE patients only will have a series of follow up appointments scheduled after the 1 month follow up appointment to adjust the lens. **Your 1st adjustment to your lens is scheduled for**

on _____ at _____ in the PORTSMOUTH SOMERSWORTH OFFICE*

***NOTE: All LAL adjustments require dilation. These appointments are generally 1 - 1 ½ hours**

PREPARING FOR YOUR CATARACT EVALUATION

By now, you've met with a counselor for a pre-evaluation to discuss lens options. Our premium lenses may significantly reduce or eliminate your need for glasses, which will be further addressed during your full evaluation with the doctor. They will review test results from your pre-evaluation, which include:

- **Biometry Tests:** To calculate the power of the artificial lens that will be implanted.
- **Optical Coherence Tomography (OCT):** To assess the health of your macula.
- **Corneal Topography:** For additional measurements if you're considering premium lenses, have had LASIK, or have significant astigmatism.

WHAT TO BRING TO YOUR CATARACT EVALUATION:

- **This Book with Your Consent Forms:** Pages 16, 17 and the Informed consent pages in the back.
- **Medication List:** Include all prescriptions, over-the-counter meds, vitamins, and supplements.
- **Specialist Contact Info:** If you currently see a specialist for any medical conditions, such as a cardiologist, it is important that you bring their contact information in case we need to have a better understanding of any of your current medications or health conditions.
- **List of Concerns:** Note any vision or health issues, even if they seem unrelated.
- **Support Person:** Having someone with you can help remember important information.
- **Transportation:** If unsure about driving post-dilation, plan for a ride. Dilation can cause light sensitivity and blurred near vision.
- **Eye Drops:** Use artificial tears if you have dry eyes to improve testing outcomes.
- **Documentation:** Bring any necessary health proxy, power of attorney, or translator documents.

DURING YOUR 60-90 MINUTE CATARACT EVALUATION, EXPECT:

- A review of your medical history and any family eye conditions.
- A vision test, including reading a chart and a slit lamp exam.
- Additional tests if necessary
- Dilation eye drops will be administered.
- Cornea (outside surface of eye) measurements will be taken. To ensure accurate measurements, please do not wear contact lenses prior to this appointment. Wear glasses only:
 - Soft lenses: remove 5 days prior.
 - Toric lenses: remove 14 days prior.
 - Hard lenses: remove 3 weeks prior.

At the end of your evaluation, you may be meeting with a surgical coordinator. If you do not meet with a coordinator, expect a call within two weeks to discuss your surgery options and the possibility of scheduling surgery date(s) with follow-up appointments.

SURGERY CHECKLIST

SCHEDULING SURGERY

- Make sure you know the date(s) for your surgery.**
 - If you are only having one eye treated, you should have 1 copy of this book.
 - If you are having 2 eyes treated, you should have 2 copies of this book.
- Make sure you know the date / time location of your follow-up appointments. You will have 1 or 2 follow-up appointments already scheduled after EACH surgery date.**
 - 1st post-op appointment – either the same day of surgery or the day after.
 - This appointment will be with your surgeon in one of our Eyesight offices.
 - 2nd post-op appointment – 2-5 weeks following surgery.
 - This appointment will be with a doctor in one of our Eyesight offices.
- Make sure you have turned in all your consent forms. We cannot schedule your surgery until you have signed:**
 - “Informed Consent for Cataract Surgery”.
 - “Lens Choice and Informed Consent for Cataract Surgery” (this was the form you signed with your surgeon at your Cataract Evaluation).
 - Additional forms, such as Health Plan Denials and Personal Obligation / Cash Pay (pg. 17), may be required if you do not have insurance or Authorization to Perform Services (pg. 16) for premium packages.
- If you have a Health Proxy, DPOA (dual power of attorney), require a translator, or need additional support for surgery, please make sure we have this information on file.**
- If you have a cardiologist or a specialist who needs to approve your surgery, please make sure we have their contact information and/or written documentation authorizing you to proceed with surgery.**

1- 2 WEEKS PRIOR TO SURGERY

- Arrange to pay for any out-of-pocket payments that may be necessary (premium lens packages).**
 - Eyesight will collect physician fees.
 - The surgical center will collect facility and lens fees.
- Make sure you have your pre/post operative eyedrops.**
 - Imprimis (available at any Eyesight location) or 3 prescriptions (called into your pharmacy).
 - Klarity-C compounded lubricating drops or lubricating over the counter eyedrops.
 - Any ocular ointments that may have been called in (usually for glaucoma or retina combined procedures).
- 1 week prior to surgery, stop taking Ozempic or other weight loss semaglutides.**
- Be prepared to receive a call from a nurse at the surgical center.**
 - They will discuss your medical history as well as the current prescriptions and medications you are using.
 - They will review pre and post-surgery information.

1-2 BUSINESS DAYS PRIOR TO SURGERY

- You will receive a call from the surgical center with your arrival time.**

DAY BEFORE SURGERY

- Review page 4-6 of this book for complete instructions.**
- Do not eat anything after midnight or your surgery will be postponed.**
- Be prepared to use your eyedrops in the morning (see page 9 or 10 of this book).**

Thank you for trusting Eyesight Ophthalmic Services for your eyecare needs.

PRIOR TO CATARACT SURGERY

PRE-SURGICAL EYE DROPS – After cataract surgery, patients are often instructed to instill multiple prescription drops to prevent infection, inflammation, and pain, and each drop is on a different schedule. This can be confusing, so we have compounded your multiple eyedrops into one prescription called Imprimis. It contains all the prescriptions you need! Patients who choose an upgraded lens package will receive eyedrops at no additional charge. **Basic Lens patients will need to purchase their Imprimis eyedrops prior to surgery. NOTE: All prescriptions or Imprimis eyedrops will need to be picked up PRIOR to surgery.** They are available at all Eyesight locations during normal business hours.

START THE IMPRIMIS DROPS ON: _____ in _____ **eye only.**

- **Beginning 1 hour BEFORE YOU LEAVE HOME**, (regardless of how long it takes you to get to the surgery center) use the IMPRIMIS eyedrops every 15 minutes for a total of 4 times. i.e., if you leave home at 9:00am, you would use the IMPRIMIS eyedrops at 8:00, 8:15, 8:30 and 8:45.
Note: Shake eye drop bottle well prior to use. Note that it is only necessary to use **1 drop at a time** from the bottle *regardless of what the package insert says.*
- Bring your eye drops and Surgery Drop Schedule to all follow-up appointments.
- **Please follow the eye drop chart you were instructed to use in this book on page 9 or 10.**

If you are using prescribed eyedrops (NOT IMPRIMIS eyedrops), please follow the eyedrop instructions on page 10 of this document.

If you regularly use other prescription eyedrops, please consult with your surgeon regarding their use before or after, surgery. If you use Xiidra, Restasis or Cequa, use 1 drop the morning of surgery, and then resume 1 WEEK after surgery,

SURGERY PREPARATION

- **You must have** responsible adult designated to accompany you to and from surgery. You will be asked to identify this person prior to surgery.
- For optimal results with your Optiwave Enhanced Treatment, Toric Lens Implant, or Presbyopic Lens Implant, **refrain from wearing contact lenses for 7 days prior to surgery, 3 weeks for rigid/GP lenses.**
- **On the night before surgery, or the morning of, take a bath or shower and wash your hair thoroughly.** In the morning, wash your face with soap and water. Please make sure you remove all mascara or eyeliner.
- **DO NOT EAT OR DRINK ANYTHING AFTER MIDNIGHT THE NIGHT BEFORE SURGERY! This includes coffee, toast, juice, etc.**
- **Take your usual morning medications with a sip of water.** If you take **INSULIN or DIABETIC medication**, the intake nurse from the surgery center will provide instructions on how and when to take your medication. As always, bring your insulin with you to surgery.
- If you use **Ozempic or weight loss semaglutides**, please stop using them 1 week before surgery.
- **BRING SUNGLASSES, GLASSES (if you wear prescription lenses), YOUR SURGERY BAG, AND EYE DROPS WITH YOU TO SURGERY.**
- **Wear loose-fitting clothing, slip on shoes (no boots), and a BUTTON-DOWN SHIRT to your surgery.** Do not wear makeup, ANY jewelry, nail polish, hairspray, perfume/cologne, or lotions. Deodorant is fine.
- **There are usually no medication restrictions for cataract surgery.** You may continue taking aspirin products, Coumadin, Plavix, etc., unless otherwise directed by your surgeon.

DURING CATARACT SURGERY

You must have a responsible adult designated to accompany you to and from surgery. You will be at the surgery center for 1-2 hours.

After check-in, you will then be brought to the “short stay” area of the operating room. Many people, (doctors and nurses alike) will speak with you, and several consent forms must be signed.

An intravenous line will be placed in your arm and several eye drops will be placed in your eye(s). You will then be brought to the operating room, and you will be given oxygen and given medication through an intravenous line to relax you.

You will be partially sedated during the procedure. The eye and skin around the eye will be cleaned and a drape will be placed over your body exposing only the operated eye. The actual procedure is usually brief – roughly 20 minutes. Recovery is fairly quick and, once cleared by the medical team, you will be ready to go home and rest.

You will have an appointment shortly after surgery and then a few weeks to a month later. You will need to pick up your prescriptions at your pharmacy or IMPRIMIS eyedrops at our office prior to surgery. You will take these drops in the operated eye for 2-6 weeks depending on your recovery.

AFTER CATARACT SURGERY

1. We prefer that you refrain from most activities for the rest of the day, and that someone stays with you until the day after surgery. Your eye may be slightly sore, itchy, scratchy, or feel a little watery.
2. Your vision may be blurry the day of surgery, and it may take several days for your vision to clear. If your eye is patched, we prefer that you keep the eye patch on, unless otherwise directed.
3. If you have both eyes done within a short period of time, you may notice one eye healing more quickly and seeing better at a faster rate. Each surgery may seem like a different experience. This is not unusual. You may also need glasses to improve your vision for distance as well as up-close.
4. Eye protection is recommended outdoors (sunglasses or glasses) and a shield at bedtime for **1 week**. Use your own discretion when indoors. Your surgeon assures you that no harm will come to your eyes if you choose to wear your “old” glasses. **Note: Light Adjustable Lens (LAL) patients MUST wear sunglasses outdoors AT ALL TIMES (or where UV exposure is possible) until your lens is locked.**

Daytime – regular glasses or sunglasses

Bedtime/Naps – Eye Shield

When putting on your eye shield, place the “arm” above your nose and place a piece of tape above and below so you can still see through it.

If you are provided with an eyepatch instead of a shield, the eyepatch is not clear to see through, but should be taped on the face in the same manner.



Eye patch



Eye shield

No restrictions on bathing, showering, shampooing hair, having a permanent, or drying your hair, BUT **NO swimming** underwater for 3 weeks. **No eye makeup** for one week.

5. **YOU MAY** travel in an airplane, read, and watch T.V.
6. **YOU MAY** bend over to put on shoes, socks, or pick up things (NOT OVER 35lbs.)
7. **YOU MAY** sleep in any position.
8. Occasionally, your vision may be good enough to drive as soon as the day after surgery. Ask your surgeon if you are not sure. Please be aware that your eye may stay dilated for 1 to 2 days after surgery.
9. Please refrain from any vigorous physical activity that might increase your chances of falling and hitting your “operated” eye (such as skiing, skating, tennis, etc.) for **1 WEEK**. It is okay for you to take a walk or play golf as long as you are wearing eye protection. If you are unsure, please ask.
10. It is not unusual to experience watering, a foreign body or a scratchy sensation for the first few weeks after surgery as the eye heals. It will improve with time.
11. While it is okay to use lubricant eye drops, i.e. “artificial tears”, please do not begin using them until at least 1 week after surgery. **If you had a premium lens implant or Optiwave**, you were provided with a compounded lubricating drop called Klarity-C. This should be used TWICE A DAY (am and pm), beginning 1 WEEK AFTER SURGERY. Use the Klarity-C until the bottles are gone.

PLEASE CALL THE OFFICE AS SOON AS POSSIBLE if you experience sudden intense pain or a dramatic change of vision in the “operated” eye.

MEDICATIONS

- Most patients will receive Imprimis eye drops, which contain Prednisolone Acetate, Moxifloxacin, and Nepafenac. These can be picked up at any Eyesight office during regular hours.
- For **Optiwave** or **Premium Lens** procedures, you’ll receive both Imprimis and Klarity-C drops (one bottle per eye) after your payment is processed. If additional prescriptions, such as ointments, are required, they will be called into your pharmacy.
- For **Basic Lens** procedures, you’ll need to purchase one bottle of Imprimis per eye or fill your prescribed eyedrops. A majority of patients will have increased symptoms of dry eye after surgery, so we also recommend over the counter lubricating eyedrops or Klarity-C lubricating drops starting one week after your cataract surgery, especially if you're having surgery on both eyes.



Note: If you're allergic to any Imprimis ingredients, we'll send a different prescription to your pharmacy.

- **Post-Surgery Eye Drop Instructions** - After surgery, use Imprimis in the operated eye four times a day (e.g., 8 am, 12 pm, 4 pm, and 8 pm) for 14 days. If advised by your doctor, reduce it to twice daily (morning and night). You'll likely need to use the drops for at least four weeks, but your doctor will inform you of any adjustments. One bottle per eye should suffice, and your surgeon will let you know if you need more.
- **Reminder:** Bring all your eyedrops, ocular ointments (if prescribed) and surgery instructions to your surgery and all follow-up appointments.
- **Refilling Prescriptions** - If you run out of Imprimis, or if you would like to purchase more Klarity-C, you can get more at any Eyesight location. If you need to refill your prescriptions at the pharmacy, a refill authorization was already sent.

If you regularly use other prescription eyedrops, such as Restasis, Xiidra or Cequa, please consult with your surgeon regarding their use before or after surgery.

Your family of Eyesight staff is here to assist you with every aspect of caring for your eyes.




Lucian Szmyd, MD



Kinley Beck, MD




Christopher Turner, OD



Warren Goldblatt, MD



Jennifer Ling, MD



Lauren McLoughlin, OD



N. Timothy Peters, MD



Jason Szelog, MD



Janet Rand, OD



Marsha Kavanagh, MD



Nathaniel Sears, MD



Renee Lynch, OD



Timothy Sullivan, MD



Dana Graichen, MD



Hilary Hamer, OD



Claudia Bartolini, MD



Dwight Arvidson, OD



PORTSMOUTH

155 Borthwick Avenue, Suite 200 East - Portsmouth, NH 03801
Tel: (603) 436-1773 Fax: (603) 427-0655

SOMERSWORTH

267 Route 108 - Somersworth, NH 03878
Tel: (603) 692-7500 Fax: (603) 692-7575

EXETER

McReel Building-192 Water Street - Exeter, NH 03833
Tel: (603) 778-1133 Fax: (603) 778-1055

KITTERY, ME

99 US-1, Suite B - Kittery, ME 03904
Tel: (207) 439-4958 Fax: (207) 439-4313

SANFORD, ME

272 Cottage Street - Sanford, ME 04073
Tel: (207) 324-3380 Fax: (207) 490-9174